

JOURNAL OF ALGEBRA 92, 116–127 (1985)

## Twisted Polynomial Rings Satisfying a Polynomial Identity

ROBERT F. DAMIANO AND JAY SHAPIRO

Department of Mathematical Sciences, George Mason University,  
Fairfax, Virginia 22030

Communicated by E. Kleinfeld

Received September 3, 1982

### INTRODUCTION

A prime ring is called *right bounded* if every essential right ideal contains a non-zero two-sided ideal. A ring is called *fully bounded* if modulo every prime ideal the ring is right and left bounded.

It is known that even if  $R$  is a fully bounded noetherian ring (*FBN ring*) the polynomial ring  $R[x]$  may fail to be an FBN ring (for an example, see [6, p. 175]). Cauchon [1] has given necessary and sufficient conditions for  $R$  so that  $R[x]$  is FBN. In this paper we study this and related questions concerning a generalized version of the polynomial ring, namely, the twisted polynomial ring. All rings are assumed to be associative with an identity. The center of a ring  $R$  is denoted by  $Z(R)$ .

Let  $\sigma$  be a unital ring endomorphism of  $R$ . The (*right*) *twisted polynomial ring*  $R[x, \sigma]$  consists of polynomials of the form  $\sum_{i=0}^n x^i a_i$  where  $a_i \in R$ . Multiplication is defined by requiring associativity, distributivity and the relation  $ax = x\sigma(a)$ . If we assume  $R$  is right (left) noetherian, then it can be shown that  $\sigma$  an automorphism implies  $R[x, \sigma]$  is right (left) noetherian. However, for an arbitrary endomorphism  $\sigma$ ,  $R[x, \sigma]$  may fail to be noetherian on either side (for instance, see [4]). We first examine necessary and sufficient conditions for a twisted polynomial ring to be FBN. Although we are unable to completely classify when  $R[x, \sigma]$  is FBN, we are able to obtain a useful classification for an important subfamily of FBN rings, namely, noetherian PI rings. In particular, we determine precisely when  $R[x, \sigma]$  is a noetherian PI ring in case  $\sigma$  is an automorphism (see Corollary 10). In Section 2, we discuss conditions under which twisted polynomial rings have “enough” localizable sets of prime ideals in the sense of Müeller [12].

## 1. FULLY BOUNDED NOETHERIAN TWISTED POLYNOMIAL RINGS

In this section, both necessary and sufficient conditions as to whether a twisted polynomial ring is FBN or satisfies a polynomial identity (PI) are presented. We first state the following proposition.

**PROPOSITION 1.** *Let  $S$  be a unital subring of a ring  $T$  such that  ${}_S T$  is a finitely generated free left  $S$ -module and let  $H = \text{End}({}_S T)$ . If there exists a basis for  $T$  over  $S$  containing the identity element  $1_T$ , then  $T$  is a unital subring of  $H$  such that  $H_T$  is a finitely generated free right  $T$ -module with a basis containing  $1_T$ .*

*Proof.* Since  $H = \text{End}({}_S T) = \text{Hom}_S({}_S T, {}_S T)$ , we have a natural right  $T$ -module structure on  $H$ . If  $\varphi \in H$ ,  $t \in T$ , then  $\varphi \cdot t$  is defined by the following relation:

$$(\varphi \cdot t)(x) = \varphi(x) t \quad \forall x \in T.$$

Thus, we can define a unital ring monomorphism as follows.

$$\theta: T \rightarrow H$$

$$\theta: t \mapsto 1_H \cdot t.$$

We remark that through this ring monomorphism, we can identify  $1_T$  and  $1_H$ .

Let  $\{1_T = a_0, a_1, \dots, a_n\}$  be a basis for  ${}_S T$ . Let  $\psi_i \in H$ ,  $i = 0, 1, \dots, n$ , be defined by

$$\psi_0 = 1_T$$

and

$$\psi_i(a_j) = \delta_{ij}, \quad i = 1, \dots, n$$

where  $\delta_{ij} = 1_T$  if  $i = j$  and 0 otherwise. We claim that  $\{1_T = \psi_0, \psi_1, \dots, \psi_n\}$  is a basis for  $H_T$ .

If  $\varphi \in H$ , then it can be checked that  $\varphi = \sum_{i=0}^n \psi_i t_i$  where  $t_0 = \varphi(a_0) = \varphi(1_T)$  and  $t_i = \varphi(a_i) - a_i t_0$ ,  $i = 1, \dots, n$ . If  $\sum_{i=0}^n \psi_i t_i = 0$  for some  $t_i \in T$ ,  $i = 0, 1, \dots, n$ , then  $t_j = 0$ ,  $j = 0, 1, \dots, n$ , since  $0 = \sum_{i=0}^n (\psi_i \cdot t_i)(a_j) = t_j$ . ■

If we restrict this proposition to the case where  $T = R[x, \sigma]$  is a twisted polynomial ring, we obtain the following useful corollary.

**COROLLARY 2.** *If  $\sigma$  is an automorphism of the ring  $R$ , then the following are equivalent:*

- (i)  $R[x, \sigma]$  is an FBN ring.
- (ii)  $R[x, \sigma^n]$  is an FBN ring for some integer  $n$ .

*Proof.* (ii)  $\Rightarrow$  (i) Let  $T = R[x, \sigma]$  and  $S = R[x, \sigma^n]$  for some integer  $n$ . There is an embedding from  $S$  to  $T$  given by  $x \rightarrow x^n$ . The rings  $T$  and  $S$  satisfy the hypotheses of Proposition 1. In particular, the elements  $\{1_R = 1_T, \dots, x^{n-1}\}$  form a basis of  $T$  as a right and left  $S$ -module.

We need to know that, in general, the matrix ring over an FBN ring is again FBN. To that end, let  $A$  be an arbitrary FBN ring and let  $B = M_n(A)$ , the ring of  $n \times n$  matrices over  $A$ . Clearly,  $B$  is noetherian. We might as well assume  $A$  is prime, so  $B$  is also prime. Thus  $B$  has a simple classical quotient ring  $M_n(Q)$  where  $Q$  is the quotient ring of  $A$ . Let  $I$  be an essential right ideal of  $B$ . Then  $I$  must contain a regular element  $C$  which is invertible in  $M_n(Q)$  by a matrix  $D$ . The elements of  $D$  are of the form  $a_{ij}q_{ij}^{-1}$ , but by common denominators the  $q_{ij}$  may be replaced by a single  $q$ . So  $CD = 1$ ,  $C(Dq) = 1q$  and  $Dq \in B$ . Thus  $I \cap A$  contains the regular element  $q$ . Since  $A$  is FBN,  $I \cap A$  contains a non-zero two-sided ideal  $J$ . Observe that  $JB = M_n(J)$  is a two-sided ideal of  $B$  contained in  $I$  and so  $B$  is fully bounded.

Returning to our particular case we see that  $H = \text{End}({}_S T) \simeq \text{End}(T_S)$  is an FBN ring. By [1, Proposition 2.1]  $T$  is an FBN ring. ■

(i)  $\Rightarrow$  (ii) This follows immediately from [1, Proposition 2.1]. ■

Later in this section, we examine a subfamily of FBN rings, namely, noetherian PI rings. For these rings the following corollary to Proposition 1 is useful. Let  $S$  and  $T$  be defined as in Proposition 1.

**COROLLARY 3.**  *$S$  satisfies a polynomial identity if and only if  $T$  satisfies a polynomial identity.*

*Proof.* The result follows immediately from the fact that a subring of a PI ring and an  $n \times n$  matrix ring over a PI ring are themselves PI rings. ■

We use these results to attain necessary criteria for  $R[x, \sigma]$  to be an FBN ring. Let  $R$  be any noetherian ring and let  $\sigma$  be an automorphism of  $R$ . If  $N$  is the nilradical of  $R$ , one can easily check that  $\sigma(N) = N$ . Therefore there is an induced automorphism

$$\bar{\sigma}: R/N \rightarrow R/N.$$

Moreover, if  $Z(R/N)$  is the center of  $R/N$ ,  $\bar{\sigma}$  can be restricted to this ring. We denote this restriction by  $\bar{\sigma}_{Z(R/N)}$ .

**PROPOSITION 4.** *Let  $R$  be a noetherian ring with  $\sigma$  an automorphism on  $R$ . If  $R[x, \sigma]$  is an FBN ring, then there exists an integer  $m$  such that  $\bar{\sigma}_{Z(R/N)}^m = 1_{Z(R/N)}$ .*

*Proof.* First, we assume  $R$  is a prime noetherian ring. It follows easily that  $R[x, \sigma]$  is also a prime ring. Note that the monic polynomial  $x + 1$  is a

regular element of  $R[x, \sigma]$ . Regular elements in a prime noetherian ring generate essential right ideals. Hence, since  $R[x, \sigma]$  is FBN, the right ideal generated by  $x + 1$  must contain a non-zero two-sided ideal  $I$ . Let  $f(x) = x^n a_n + x^{n-1} a_{n-1} + \dots + a_0$  be a non-zero element of minimal degree in  $I$ . Since  $x + 1$  divides  $f(x)$ , we know that at least one of the non-leading coefficients of  $f(x)$ , say,  $a_i$ ,  $i \neq n$ , is non-zero. If  $b$  is any element of  $Z(R)$ , then  $g(x) = b \cdot f(x) - f(x) \cdot \sigma^n(b)$  is an element of  $I$ . Since  $\sigma^n(b)$  is a central element  $g(x)$  has degree less than  $f(x)$ . Thus  $g(x) = 0$ . In particular, since  $a_i \neq 0$  and since  $\sigma^i(b) - \sigma^n(b)$  is in the center of the prime ring  $R$ , it follows that  $\sigma^i(b) = \sigma^n(b)$ . Therefore, if  $m = n - i$ , we see that  $\sigma_{Z(R)}^m = 1_{Z(R)}$ .

Now let  $R$  be a semiprime noetherian ring. By Goldie's theorem [15, II, 2.4] there exists a finite number of minimal prime ideals  $P_1, \dots, P_k$  of  $R$ . Since  $\sigma$  is an automorphism,  $\sigma(P_i) = P_j$  for some  $j$ . Since the  $\sigma$  acts as a permutation on the set  $\{1, \dots, k\}$ , there exists an integer  $n$  such that  $\sigma^n(P_i) = P_i$  for all  $i$ . If  $\rho = \sigma^n$ , then by Corollary 2 we know that  $R[x, \rho]$  is FBN. Therefore  $R[x, \rho]/P_i R[x, \rho] \cong R/P_i[x, \bar{\rho}]$  is a prime FBN ring. Hence, there exist an  $m_i$  such that  $\bar{\sigma}_{Z(R/P_i)}^{m_i} = 1_{Z(R/P_i)}$ . Let  $m$  equal the least common multiple of the set  $\{nm_i \mid i = 1, \dots, k\}$ . It is clear that  $\sigma^m(P_i) = P_i$  and  $\bar{\sigma}_{Z(R/P_i)}^m = 1_{Z(R/P_i)}$  for each  $i = 1, \dots, k$ . If  $b \in Z(R)$ , then the image of  $b$  is central in each  $R/P_i$ . Therefore  $b - \sigma^m(b) \in P_i$  for each  $i$ . Since  $P_1 \cap \dots \cap P_k = 0$ ,  $\sigma^m(b) = b$  and we are done.

If  $R$  is any noetherian ring with nilradical  $N$ , then the proposition follows since  $R[x, \sigma]/N[x, \sigma] \cong R/N[x, \bar{\sigma}]$ . ■

We remark that although we required a noetherian hypothesis, similar results are attainable if the ring has a finite set of minimal prime ideals with nilpotent nilradical. In the case of a ring satisfying a polynomial identity, we can prove a stronger result. We recall that given a semiprime ideal  $N$  of  $R$ , then  $\mathcal{C}(N)$  will denote the elements of  $R$  which are regular modulo  $N$ .

**LEMMA 5.** *If  $R$  is a semiprime, right Goldie ring and if  $\sigma$  is a monic endomorphism of  $R$  such that  $\sigma(\mathcal{C}(0)) \subseteq \mathcal{C}(0)$ , then for each minimal prime ideal  $P$  of  $R$ , there exists an integer  $n$  such that  $\sigma^n(P) \subseteq P$ .*

*Proof.* We first consider the case when  $R$  is a semisimple artinian ring. Let  $P_1, P_2, \dots, P_k$  be the prime ideals of  $R$  and let  $\{e_1, e_2, \dots, e_k\}$  be a complete set of orthogonal primitive idempotents of  $R$ . Therefore  $\{\sigma(e_1), \dots, \sigma(e_n)\}$  is also a complete set of orthogonal primitive idempotents. By [9, p. 54, Theorem 3], there exists a unit  $u \in R$  such that  $e_i = u^{-1} \sigma(e_i) u$ . Let  $\pi^u$  be the automorphism of  $R$  that is defined via conjugation by  $u$ . Thus the endomorphism  $\pi^u \circ \sigma$  fixes each  $e_i$ . Since any ideal of  $R$  is generated by a subset of  $\{e_1, \dots, e_n\}$  as a right ideal, it follows that  $\pi^u \circ \sigma(P_i) \subseteq P_i$ . Since  $\pi^{u^{-1}}$  is an automorphism of  $R$ , we know that  $\pi^{u^{-1}}(P_i) = P_j$  for some  $j$ . Therefore,  $\sigma(P_i) = \pi^{u^{-1}} \circ \pi^u \circ \sigma(P_i) \subseteq \pi^{u^{-1}}(P_i) = P_j$ .

Since  $\pi^u \circ \sigma$  fixes each  $e_i$ ,  $\pi^u \circ \sigma(P_i)R = P_i$ . Thus, if  $i \neq k$ ,  $\pi^u \circ \sigma(P_k) \not\subseteq P_i$  and it follows that  $\sigma(P_i)$  and  $\sigma(P_k)$  are not contained in the same  $P_j$  for  $i$  unequal to  $k$ . By a counting argument, we see that for each  $j$ ,  $P_j$  contains  $\sigma(P_i)$  for one and only one  $i$ . More generally, for each integer  $m$ ,  $P_j$  contains  $\sigma^m(P_i)$  for one and only one  $i$ . Clearly for each prime ideal  $P$  there exists distinct integers  $m$  and  $t$  such that  $\sigma^m(P) = \sigma^t(P)$  are contained in the same prime ideal. Thus  $\sigma^{m-1}(P)$  and  $\sigma^{t-1}(P)$  also must be contained in the same prime ideal. By continuing inductively in this manner, we have the special case.

Suppose  $R$  is a semiprime Goldie ring with endomorphism  $\sigma$ . It follows that the classical ring of quotients  $Q$  is a semisimple ring with prime ideals  $\bar{P}_1, \dots, \bar{P}_k$  which are the localizations of the minimal prime ideals  $P_1, \dots, P_k$  of  $R$ . Moreover, since  $\sigma$  takes regular elements to regular elements it can be extended to a monic endomorphism of  $Q$ . The result now follows. ■

We now derive a stronger version of Proposition 4 in the case of PI ring.

**PROPOSITION 6.** *Let  $R$  be a semiprime, right Goldie ring with  $\sigma$  a monic endomorphism such that  $\sigma(\mathcal{C}(0)) \subseteq \mathcal{C}(0)$ . If  $R[x, \sigma]$  is a PI ring, then an integer  $n$  exists such that  $\sigma^n_{Z(R)} = 1_{Z(R)}$ .*

*Proof.* Assume  $R$  is a prime ring. It follows that  $R[x, \sigma]$  is a prime ring and thus the two-sided ideal  $xR[x, \sigma]$  contains a non-zero central element  $f(x)$  [13, Theorem 1.7.11]. We note that  $f(x)$  must be of degree at least one. Let  $f(x) = x^n a_n + x^{n-1} a_{n-1} + \dots + x a_1$ . Since  $f(x)$  is central, it follows that each monomial  $x^i a_i$  is central. In particular  $x^n a_n$  is central. If  $b \in Z(R)$ , then  $b x^n a_n = x^n a_n b$ . This implies  $\sigma^n(b) a_n = b a_n$  and hence  $(\sigma^n(b) - b) a_n = 0$ . However, since  $x^n a_n$  is a central element of a prime ring it must be a regular element of that ring. It follows that  $a_n$  is a regular element of  $R$ . Thus  $\sigma^n(b) = b$ .

Let  $R$  be semiprime. By Lemma 5, there exists an integer  $n$  such that for every minimal prime ideal  $P$  of  $R$ ,  $\sigma^n(P) \subseteq P$ . Thus  $\hat{P} = \sum x^i P$  is an ideal of  $R[x, \sigma^n]$ . If  $\bar{\sigma}^n$  is the induced endomorphism of  $R/P$ , then  $R[x, \sigma^n]/\hat{P} \cong R/P[x, \bar{\sigma}^n]$ . The ring  $R/P[x, \bar{\sigma}^n]$  is a PI ring. We claim  $\bar{\sigma}^n$  is a monomorphism. Suppose  $\bar{\sigma}^n(P) \supset P$ , then  $\bar{\sigma}^n(P)$  must contain a regular element. Since  $\sigma$  takes regular elements to regular elements, so does  $\sigma^n$ . Hence  $P$  must contain a regular element. However,  $P$  is a minimal prime ideal of a semiprime, Goldie ring, so it is the annihilator of an ideal of  $R$ , a contradiction. Thus  $\bar{\sigma}^n(P) = P$  and  $\bar{\sigma}^n$  is a monomorphism. Hence,  $R/P[x, \bar{\sigma}^n]$  is a prime ring. We have reduced the problem to that of a prime PI ring and the result follows as in Proposition 4. ■

So far, we have produced necessary conditions for a twisted polynomial ring to be FBN or to satisfy a polynomial identity. In Propositions 7, 8 and

9 we give sufficient conditions for  $R[x, \sigma]$  to be FBN or to satisfy a polynomial identity. While we do not have a complete classification in the FBN case, we do have the following result in that direction.

**PROPOSITION 7.** *Let  $R$  be an FBN ring with nilradical  $N$  such that  $R[x]$  is an FBN ring. If  $\sigma$  is an automorphism of  $R$  such that  $\bar{\sigma}^n = 1_{R/N}$  for some integer  $n$ , then  $R[x, \sigma]$  is an FBN ring.*

*Proof.* Since  $N[x, \sigma]$  is nilpotent and hence contained in every prime ideal, it suffices to prove that  $R[x, \sigma]/N[x, \sigma] \cong R/N[x, \bar{\sigma}]$  is an FBN ring. But  $R/N[x, \bar{\sigma}^n] \cong R/N[x]$  is an FBN ring. By Corollary 2,  $R/N[x, \bar{\sigma}]$  is an FBN ring. ■

For a complete classification of when  $R[x]$  is an FBN ring, we refer the reader to a paper of Cauchon [1]. Recall that an important class of FBN rings are those noetherian rings algebraic over their center. If  $R$  is such a ring, then so is  $R[x]$ . Hence, Proposition 7 is applicable in this case. For these rings, we can attain a result somewhat more general than Proposition 7.

**PROPOSITION 8.** *Let  $R$  be a noetherian ring algebraic over its center with nilradical  $N$ . If  $\sigma$  is an automorphism of  $R$  such that the induced map  $\bar{\sigma}^n$  is conjugation by some unit  $\bar{u}$  in  $R/N$ , then  $R[x, \sigma]$  is an FBN ring.*

*Proof.* As in Proposition 7, we can assume  $R$  is a semiprime ring and that  $\sigma$  is conjugation by a unit  $u$  of  $R$ . Since  $\sigma$  is an automorphism,  $R[x, \sigma]$  is noetherian [11]. We first claim that  $R[x, \sigma]$  is algebraic over its center. We note that a monomial of the form  $x^m u^{-m} a$  with  $a \in Z(R)$  is central in  $R[x, \sigma]$ . Clearly the elements of  $R$  are algebraic over the center of  $R[x, \sigma]$ . Thus we will be done if we can show  $x$  satisfies a polynomial of the form  $y^n a_0 + y^{n-1} x u^{-1} a_1 + y^{n-2} x^2 u^{-2} a_2 + \cdots + x^n u^{-n} a_n$  where each  $a_i \in Z(R)$ . If we substitute  $x$  for  $y$  in this equation, we obtain the relation

$$x^n(a_0 + u^{-1}a_1 + \cdots + u^{-n}a_n) = 0.$$

Thus, it suffices to show that there exist elements  $a_i \in Z(R)$  such that  $a_0 + u^{-1}a_1 + \cdots + u^{-n}a_n = 0$ . However, if we multiply this equation by  $u^n$ , we see that it suffices to show  $u$  satisfies a polynomial over  $Z(R)$ . Since  $R$  is algebraic over  $Z(R)$  we have the claim. Finally, we want to show that if a ring  $T$  is a noetherian ring algebraic over its center  $Z$ , then it is an FBN ring. Without loss of generality assume  $T$  is prime and algebraic over  $Z$ . Let  $I$  be an essential right ideal of  $R$ , we must show that  $I$  contains a non-zero two-sided ideal. By Goldie's Theorem,  $I$  contains a regular element  $c$ . Let  $p(x) = x^n a_n + x^{n-1} a_{n-1} + \cdots + x a_1 + a_0$  be the polynomial over  $Z$  of least degree satisfied by  $c$ . If  $a_0 \neq 0$  then  $a_0 \in I \cap Z$  so  $I$  contains a two-sided ideal. If

$a_0 = 0$ , then  $c(c^{n-1}a_n + c^{n-2}a_{n-1} + \cdots + a_1) = 0$  and since  $c$  is regular,  $c$  satisfies a polynomial of degree smaller than  $p(x)$ , a contradiction. Thus  $R[x, \sigma]$  is FBN. ■

We now give sufficient conditions for  $R[x, \sigma]$  to satisfy a polynomial identity.

**PROPOSITION 9.** *Let  $R$  be a PI ring with nilpotent nilradical  $N$  such that  $N$  is the intersection of finite number of prime ideals. If  $\sigma$  is a monic endomorphism of  $R$  such that  $\sigma(N) \subseteq N$ ,  $\sigma(\mathcal{C}(N)) \subseteq \mathcal{C}(N)$ , and  $\bar{\sigma}^n = 1_{Z(R/N)}$  for some integer  $n$ , then  $R[x, \sigma]$  is a PI ring.*

*Proof.* Since  $\sigma$  is a monic endomorphism, it follows that  $\sigma^{-1}(N) \subseteq N$ . Thus if  $\sigma(N) \subseteq N$ , we have an induced endomorphism  $\bar{\sigma}$  of  $R/N$  which is monic and sends regular elements to regular elements. Again, as in our previous propositions, we can now assume  $R$  is semiprime.

Since  $R$  has a finite number of minimal prime ideals,  $R$  has a semisimple artinian ring of quotients  $Q$  [13, Theorem 1.7.34]. It follows that  $Q$  is a PI ring, finite dimensional over its center  $Z(Q)$ , and  $Z(Q)$  is the classical ring of quotients of the center of  $R$  [13, Theorem 1.7.20].

Let  $\hat{\sigma}$  be the extension of  $\sigma$  to  $Q$ . We note that  $\hat{\sigma}^n = 1_{Z(Q)}$  since  $Z(Q)$  is classical ring of quotients of  $Z(R)$ . Since  $Q$  is finitely generated, free over  $Z(Q)$  with a basis containing the identity, the same can be said for  $Q[x, \hat{\sigma}^n]$  over  $Z(Q)[x]$ . By Corollary 3,  $Q[x, \hat{\sigma}^n]$  satisfying a polynomial identity implies that  $Q[x, \hat{\sigma}]$  is a PI ring. Since  $R[x, \sigma] \subseteq Q[x, \hat{\sigma}]$ , we are done. ■

**COROLLARY 10.** *If  $R$  is a noetherian PI ring with automorphism  $\sigma$  and nilradical  $N$ , then the following are equivalent:*

- (i)  $R[x, \sigma]$  is a noetherian PI ring.
- (ii)  $\bar{\sigma}^n = 1_{Z(R/N)}$  for some integer  $n$ .

*Proof.* (i)  $\Rightarrow$  (ii) If  $R[x, \sigma]$  is a noetherian PI ring, then the result follows from Proposition 4.

(ii)  $\Rightarrow$  (i) The results follows from Proposition 9. ■

We now give two examples. In the first, we have a commutative artinian ring  $R$  and an automorphism  $\sigma$  such that  $\sigma^n \neq 1_R$  for any  $n$ , but whose induced automorphism is the identity on  $R/N$ . In the second example we give an automorphism  $\sigma$  such that no power of  $\sigma$  is the identity on all of  $R$  but  $\sigma$  restricted to the center is the identity. In both cases  $R[x, \sigma]$  satisfies a polynomial identity by the results of this section.

**EXAMPLE 1.** Let  $R = \left\{ \begin{pmatrix} a & 0 \\ b & a \end{pmatrix} \mid a, b \in \mathbb{C}, \text{ the complex numbers} \right\}$ . Let  $u$  be a non-zero element of  $\mathbb{C}$  which is not a root of unity. Define

$$\sigma: R \rightarrow R$$

$$: \begin{pmatrix} a & 0 \\ b & a \end{pmatrix} \rightarrow \begin{pmatrix} a & 0 \\ ub & a \end{pmatrix}.$$

This is an automorphism, no power of which is the identity. It is clear that  $\bar{\sigma}$  is the identity on  $R/N$ .

EXAMPLE 2. Let  $R$  be the ring of  $2 \times 2$  matrices over the rationals and let  $u = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$ . Let  $\sigma$  be conjugation by  $u$ . Of course  $\sigma = 1_{Z(R)}$  and no power of  $\sigma$  is the identity.

To find a noetherian, PI ring  $R$  and an automorphism  $\sigma$  such that  $R[x, \sigma]$  is not a PI ring one need only take a commutative field  $R$  and an automorphism  $\sigma$  of infinite order.

## 2. LOCALIZATION OF TWISTED POLYNOMIAL RINGS

Recall that a finite set  $\{P_1, P_2, \dots, P_n\}$  of prime ideals is said to be *localizable* if  $\mathcal{C}(N)$  is an Ore set where  $N = \bigcap_{i=1}^n P_i$ . In an FBN ring, Müller [12] calls a minimal localizable set of prime ideals a *clan*. He showed that a prime ideal is contained in at most one clan. We will use  $[P]$  to denote the unique clan containing  $P$  (when the clan exists). If every prime of an FBN ring belongs to a localizable set we say the ring has *enough clans*. Moreover, if  $R$  is an FBN ring with enough clans such that each clan is a singleton set, we say  $R$  is *stable* [3].

From the first section we know that  $R[x, \sigma]$  will be FBN under certain circumstances. In this section we want to discuss when primes ideals of  $R[x, \sigma]$  are localizable.

PROPOSITION 11. *Let  $R$  be a simple artinian ring with automorphism  $\sigma$ . If  $R[x, \sigma]$  is FBN, then  $R[x, \sigma]$  is stable.*

*Proof.* We note that  $R[x, \sigma]$  has global dimension 1, i.e.,  $R[x, \sigma]$  is a hereditary ring [5, Corollary]. As we said before, it is easy to see that  $R[x, \sigma]$  is a prime ring. Therefore, it follows from [4, Proposition 2.2] and [2, Theorem 3.3] that we need only show  $R[x, \sigma]$  has no idempotent non-zero two-sided ideals. It is known that  $R[x, \sigma]$  is a principal right ideal ring (for instance, see [10, Theorem 3.1]). So, if  $I$  is two-sided ideal, then  $I = f \cdot R[x, \sigma]$  for some  $f \in R[x, \sigma]$ . If  $I^2 = I$ , then  $f \cdot R[x, \sigma] \subseteq fR[x, \sigma] \cdot fR[x, \sigma] \subseteq f^2 \cdot R[x, \sigma]$ . Thus  $f = f^2g$  for some  $g \in R[x, \sigma]$ . Therefore, either  $f \cdot g = 1$ , a contradiction, or  $f$  is a zero divisor. However,  $f \cdot R[x, \sigma]$  is an ideal of a prime ring, so it must contain a regular element. Hence,  $f$  itself must be regular. Thus  $I^2 \neq I$  and we are done. ■



We will need some terminology before going to the next result. The Krull dimension of a ring  $R$ , in the sense of Gordon and Robson [7], will be denoted  $K\text{-dim } R$ . If  $R$  is an FBN ring then  $K\text{-dim } R$  is the same as the classical Krull dimension of  $R$ . If  $M$  is an  $R$  module,  $E(M)$  will denote the injective full of  $M$ .

Recall that a ring  $S$  is called a finite normalizing extension of a unital subring  $R$  provided that  $S$  is finitely generated as an  $R$ -module by elements which normalize  $R$ ; that is,  $S = \sum_{i=1}^n Ra_i$  with  $Ra_i = a_iR$  for each  $i$ .

**THEOREM 12.** *Let  $S \subseteq T$  be a finite normalizing extension of  $S$  such that  $T$  is free as a left and right  $S$ -module. Suppose a normalizing basis for  $T$  over  $S$  is of the form  $\{1, b, b^2, \dots, b^{n-1}\}$  where  $b$  is regular in  $T$  with  $b^nQ = Qb^n$  for all primes  $Q$  in  $S$ . If  $S$  is FBN with enough clans, then  $T$  is FBN with enough clans.*

*Proof.* By Proposition 1 and [1, Proposition 2.11]  $T$  is an FBN ring. Let  $P$  be a prime ideal of  $T$ . By Heinicke and Robson [8, Corollary 2.12], there exist a finite number of prime ideals of  $S$ ,  $\{Q_1, \dots, Q_r\}$  minimal over  $P \cap S$ . Moreover,  $S/Q_i \cong S/Q_j$   $1 \leq i, j \leq r$ , and for each  $i$ ,  $1 \leq i \leq r$ , there exists  $j$ ,  $1 \leq j \leq n-1$ , such that  $\bar{Q}_i \bar{b}^j = \bar{b}^j \bar{Q}_1$  where  $\bar{X}$  denotes the image of the set  $X$  in  $T/P$  [8, Theorem 2.11].

Since  $b$  is regular in  $T$  and  $bS = Sb$ ,  $b$  induces a ring automorphism  $\sigma$  of  $S$  defined by  $sb = b\sigma(s)$ . Thus

$$\bar{Q}_i \bar{b}^j = \bar{b}^j \bar{Q}_1 = \overline{\sigma^j(Q_1)} \cdot \bar{b}^j.$$

If  $b \notin P$ , then  $\bar{b}$  is a normalizing element of  $T/P$ , i.e.,  $T/P \cdot \bar{b} = \bar{b} \cdot T/P$ . Hence  $\bar{b}$  is regular in  $T/P$ . Therefore,  $\bar{Q}_i = \sigma^j(Q_1)$  and since  $P \cap S \subseteq Q_i$  it follows that  $\sigma^j(Q_1) \subseteq Q_i$ . However,  $K\text{-dim}(S/\sigma^j(Q_1)) = K\text{-dim}(S/Q_1) = K\text{-dim}(S/Q_i)$ , and hence  $\sigma^j(Q_1) = Q_i$ .

If  $b \in P$ , it is clear that  $S/P \cap S = T/P$ , so  $P \cap S$  is prime. In either case, the prime ideals of  $S$  minimal over  $P \cap S$  are a subset of the set  $\{\sigma^i(Q) \mid Q \text{ is a minimal prime over } P \cap S, 0 \leq i \leq n-1\}$  (note  $\sigma^n(Q) = Q$  by hypothesis). Let

$$U_p = \{Q \mid Q \in [\sigma^j(Q_p)] \text{ where } Q_p \text{ is a prime minimal over } P \cap S\}$$

and

$$V_p = \{P' \mid P' \text{ is prime ideal of } T \text{ and there exists } Q \in U_p \text{ minimal over } P'\}.$$

We note that since  $S$  has enough clans,  $U_p$  is a finite set. By [8,

Corollary 2.12],  $V_p$  is a finite set. Let  $P' \in V_p$ . Suppose there exists a  $T$ -homomorphism

$$\varphi: E(T/P') \rightarrow E(T/P'')$$

where  $P''$  is a prime ideal of  $T$  and  $K\text{-dim}(T/P') = K\text{-dim}(T/P'')$ . In order to show  $T$  has enough clans, it suffices to prove that  $P'' \in V_p$  [12, Theorem 5].

Let  $E_S(T/P')$  be the left  $S$ -injective envelope of  $T/P'$  and consider the left  $T$ -module  $\text{Hom}_S({}_S T, E_S(T/P'))$ . By a modification of [15, Proposition 10.4],  $\text{Hom}_S(T, E_S(T/P'))$  is an injective  $T$ -module. Consider the quotient homomorphism  $\psi: T \rightarrow T/P'$ . It follows that  $P' = \text{ann}_T(\psi)$ . Therefore  $E(T/P') \subseteq \text{Hom}_S(T, E_S(T/P'))$  as a  $T$ -module. As an  $S$ -module

$$\begin{aligned} \text{Hom}_S(T, E_S(T/P')) &\cong \bigoplus_{i=0}^{n-1} \text{Hom}_S(Sb^i, E_S(T/P')) \\ &\cong \bigoplus_{i=0}^{n-1} \text{Hom}_S(Sb^i, \bigoplus_{k=1}^m E(S/N_k)) \cong \bigoplus_{k=1}^m \bigoplus_{i=0}^{n-1} \text{Hom}_S(Sb^i, E(S/N_k)) \end{aligned}$$

where  $N_k \in \{\sigma^j(Q') \mid Q' \in U_p \text{ and minimal over } P' \cap S\}$  [8, Theorem 3.6]. One checks that  $\text{Hom}_S(Sb^i, E(S/N_k))$  is isomorphic to  $E(S/M_k)$  where  $M_k \in \{\sigma^j(Q')\}$ . Thus  $E(T/P') \cong \bigoplus_{h=1}^r E(S/M_k)$  where  $M_k \in \{\sigma^j(Q')\}$ .

Since  $\varphi: E(T/P') \rightarrow E(T/P'')$  is a non-zero  $S$ -homomorphism, without loss of generality,  $\varphi(E(S/M_1)) \neq 0$ . Let  $E(S/N)$  be an  $S$ -direct summand of  $E(T/P'')$  such that there exists a non-zero  $S$ -homomorphism  $E(S/M_1) \rightarrow E(S/N)$ ,  $N$  a prime ideal of  $S$  minimal over  $P'' \cap S$ . Since  $K\text{-dim}(S/M_1) = K\text{-dim}(T/P') = K\text{-dim}(T/P'') = K\text{-dim}(S/N)$ , it follows that  $N \in \{M_1\}$ . We note that  $M_1 = \sigma^t(Q')$  where  $Q' \in U_p$  and  $0 \leq t \leq n-1$ . By [12, Proposition 2(4)], one can check  $U_p$  is closed under  $\sigma$ . Hence,  $M_1 \in U_p$  and it follows that  $N \in U_p$ . By definition,  $P'' \in V_p$ . ■

**COROLLARY 13.** *Let  $R$  be a noetherian ring integral over its center. If  $\sigma$  is an automorphism of  $R$  such that  $\sigma^n$  is conjugation by some unit of  $R$  then  $R[x, \sigma]$  is an FBN ring with enough clans.*

*Proof.* By Proposition 8  $R[x, \sigma]$  is an FBN ring. It follows from the proof of Proposition 8 that  $R[x, \sigma^n] \cong R[x^n, \sigma]$  is integral over its center. Thus  $R[x^n, \sigma]$  has enough clans [12, Proposition 9]. Clearly  $R[x^n, \sigma]$  and  $R[x, \sigma]$  satisfy the relationship of  $S$  and  $T$ , respectively, in Theorem 12. ■

**COROLLARY 14.** *Let  $R$  be a finitely generated algebra over a field. Let  $\sigma$  be an automorphism of  $R$  of finite order. If  $R$  is a noetherian PI ring with enough clans then so is  $R[x, \sigma]$ .*

*Proof.* In [14, Proposition 9] it was shown that the corollary is true if  $\sigma$  is the identity. When combined with Corollary 3 and Theorem 12 we have the desired result. ■

We note that in the preceding results there exists a non-zero integer  $j$  such that  $\sigma^j(P) = P$  for each prime  $P$  in  $R$ . In a sense this property is essential to ensure that  $R[x, \sigma]$  has enough clans.

EXAMPLE 3. Let  $R = \mathbb{C}[y]/\langle y^2 \rangle$ . We define an automorphism

$$\sigma: R \rightarrow R$$

$$\sigma: a_0 + a_1 y \rightarrow a_0 + a_1 u y$$

where  $0 \neq u \in \mathbb{C}$  such that  $u$  is not a root of unity. Consider the twisted polynomial ring  $R[x, \sigma]$ . Let  $M_c$  be the left ideal generated by  $x - c$  and  $y$ ,  $c \in \mathbb{C}$ . Note that  $y$  is a normalizing element and  $R[x, \sigma]/\langle y \rangle$  is a commutative ring. Hence  $M_c$  is a two-sided maximal ideal. We claim that if  $0 \neq a = ub$  then  $M_a$  and  $M_b$  are two maximal ideals which are linked in the sense of Müeller [12]. By [12, p. 236, Remark (2)] it suffices to show  $M_a \cap M_b \not\supseteq M_a M_b$ . Since  $y \in M_a \cap M_b$  it will suffice to show that  $y \notin M_a M_b$ .

Suppose  $y \in M_a \cdot M_b$ . Therefore

$$\begin{aligned} y &= q_1(x - a)(x - b) + q_2(x - a)y + q_3y(x - b) \\ &= q_1(x - a)(x - b) + q_2y(xu^{-1} - a) + q_3y(x - b) \\ &= q_1(x - a)(x - b) + q_2yu^{-1}(x - b) + q_3y(x - b) \\ &= q(x - b) \end{aligned}$$

where  $q_1, q_2, q_3$  and  $q \in R[x, \sigma]$ . The left side of this equation has degree 0 as a polynomial in  $x$ , while the right side has degree at least one, unless  $q = 0$ . In either case we have a contradiction. Thus  $M_b$  and  $M_{ub}$  are linked. By induction,  $M_b$  and  $M_{ukb}$  are also linked for all  $k \in \mathbb{N}$ . Since  $u$  is not a root of unity, we have infinitely many primes “linked” together. Hence  $M_b$ ,  $b \neq 0$ , does not belong to a localizable set of prime ideals [10, Theorem 5].

Given Theorem 12 and [12, Proposition 9] it is natural to ask for what PI rings  $R$  does  $R[x]$  have enough clans? Our next example, which is a variation of the previous one, shows that even if  $R$  is an artinian PI ring with only one non-zero ideal,  $R[x]$  may not have enough clans.

EXAMPLE 4. Let  $F$  be any field and let  $Q = F(z_i)$ ,  $i \in \mathbb{Z}$  be the field of rational functions of  $F$  adjoining countably many commuting indeterminates. Let  $\sigma$  be the endomorphism of  $Q$  which fixes  $F$  and sends  $z_i$  to  $z_{i+1}$ . Let

$R = Q[y, \sigma]/(y^2)$ . It is not difficult to see that  $R$  is an artinian PI ring (since modulo the nilradical  $R$  is commutative) with unique non-zero ideal  $(y)$ .

Consider the ring  $R[x]$ . For each  $z_i$  let  $M_i$  be the left ideal generated by  $x - z_i$  and  $y$ . As before  $M_i$  is a two-sided maximal ideal. We claim  $M_i$  is linked to  $M_{i+1}$ . As before it will suffice to show that  $y \notin M_i \cdot M_{i+1}$ . Assume the contrary. Then

$$\begin{aligned} y &= q_1(x - z_i)(x - z_{i+1}) + q_2(x - z_i)y + q_3y(x - z_{i+1}) \\ &= q_1(x - z_i)(x - z_{i+1}) + q_2y(x - z_{i+1}) + q_3y(x - z_{i+1}) \\ &= q(x - z_{i+1}) \end{aligned}$$

where  $q_1, q_2, q_3$  and  $q \in R[x]$ . As in the previous example we have a contradiction. So again,  $M_i$  cannot belong to a clan.

## REFERENCES

1. G. GAUCHON, Anneaux de polynomes essentiellement bornes, in "Lecture Notes in Pure and Applied Mathematics," Vol. 51, Dekker, New York, 1979.
2. A. W. CHATTERS AND A. G. HEINICKE, Localization at a torsion theory in hereditary noetherian rings, *Proc. London Math. Soc.* (3) **27** (1973), 193–204.
3. R. F. DAMIANO AND Z. PAPP, On consequences of stability, *Comm. Algebra* **9** (1981), 747–764.
4. D. EISENBUD AND J. C. ROBSON, Hereditary noetherian prime rings, *J. Algebra* **16** (1970), 86–104.
5. K. FIELDS, On the global dimension of skew polynomial rings, *J. Algebra* **13** (1969), 1–4.
6. R. GORDON, Polynomial modules over Macauley modules, *Canad. Math. Bull.* **19** (2) (1976), 173–176.
7. R. GORDON AND J. C. ROBSON, Krull dimension, *Mem. Amer. Math. Soc.* **133** (1973).
8. A. G. HEINICKE AND J. C. ROBSON, Normalizing extensions prime ideals and incomparability, *J. Algebra* **72** (1981), 237–268.
9. N. JACOBSON, Structure of rings, in "Colloquium Publications," Vol. 37, Amer. Math. Soc., Providence, R. I., 1956 (revised 1964).
10. A. V. JATEGAONKAR, Left principal ideal rings, in "Lecture Notes in Mathematics No. 123," Springer-Verlag, New York/Berlin, 1970.
11. P. MALCOLMSON AND J. SHAPIRO, The Hilbert basis theorem for skew polynomial rings, *Houston J. Math.* **8** (3) (1982), 359–368.
12. B. J. MÜELLER, Localization in fully bounded noetherian rings, *Pacific J. Math.* **67** (1976), 233–245.
13. L. H. ROWEN, "Polynomial Identities in Ring Theory," Academic Press, New York, 1980.
14. J. SHAPIRO,  $R$ -sequences in fully bounded noetherian rings, *Comm. Algebra* **7** (1979), 819–831.
15. B. STENSTRÖM, Rings of Quotients," Springer-Verlag, New York/Berlin, 1975.